

COXETER-CATALAN COMBINATORICS

- **Catalan object**: a finite set associated with the symmetric group whose cardinality equals some Catalan number
- **problem**: given an integer $m \geq 1$, parametrize a Catalan object so that the resulting set has cardinality equal to the corresponding Fuß-Catalan number, and when $m = 1$ the set coincides with the corresponding classical Catalan object
- **general problem**: recall that the symmetric group is a Coxeter group of type A ; generalize the above constructions to any finite Coxeter group

MOTIVATION

- **Tamari lattice**: a classical Catalan object that can be realized as the set of Dyck paths equipped with the rotation order
- **m -Tamari lattice**: the set of m -Dyck paths equipped with the rotation order
- **m -Tamari lattice associated with a Coxeter group W : open!**
we provide a new realization of the classical m -Tamari lattice in terms of m -tuples of Dyck paths; using the same construction (the m -cover posets), we provide an “ m -Tamari like” lattice for the dihedral group \mathfrak{D}_k


THE m -COVER POSETS

Let $\mathcal{P} = (P, \leq)$ be a bounded poset, let $\hat{0}$ denote the least element, and let $m > 0$.

- **m -cover poset $\mathcal{P}^{(m)}$** : the subposet of the direct product \mathcal{P}^m with ground set


$$\mathcal{P}^{(m)} = \left\{ (\hat{0}_1, \dots, \hat{0}_1, p_1, \dots, p_1, q_1, \dots, q_1) \mid p_1 \leq q_1, l_1 + l_2 + l_3 = m \right\}$$

EXAMPLE: THE PENTAGON LATTICE AND ITS 2-COVER POSET



THE m -DYCK PATHS AND THE m -TAMARI LATTICES


- **m -Dyck path of length $(m+1)n$** : north-east path from $(0,0)$ to (mn, n) that stays weakly above the line $x = my$
- $\mathcal{D}_n^{(m)}$ denotes the set of all m -Dyck paths of length $(m+1)n$
- $|\mathcal{D}_n^{(m)}| = \text{Cat}^{(m)}(\mathfrak{S}_n) = \frac{1}{mn+1} \binom{(m+1)n}{n}$
- **rotation order \leq_R** : covers given by exchanging an east step with the subsequent subpath



- **m -Tamari lattice**: the poset $\mathcal{T}_n^{(m)} = (\mathcal{D}_n^{(m)}, \leq_R)$

THE STRIP-DECOMPOSITION

- **strip-decomposition**: the map $\delta : \mathcal{D}_n^{(m)} \rightarrow (\mathcal{D}_n)^m$ given by “cutting the path into strips and regrouping”



THEOREM: A NEW REALIZATION OF THE m -TAMARI LATTICE

For $m, n > 0$ we have $\mathcal{T}_n^{(m)} \cong \mathbf{DM}(\mathcal{T}_n^{(m)})$, where **DM** denotes the Dedekind-MacNeille completion.

Sketch of the proof. Combine the following.

- $\mathcal{P} \cong \mathbf{DM}(\mathcal{M}(\mathcal{P}) \cup \mathcal{J}(\mathcal{P}))$, where \mathcal{P} is a finite lattice [known]
- $(\mathcal{M}(\mathcal{T}_n^{(m)}), \leq_R) \cong (\mathcal{M}(\mathcal{T}_n^{(m)}), \leq_R)$
- $(\mathcal{J}(\mathcal{T}_n^{(m)}), \leq_R) \cong (\mathcal{J}(\mathcal{T}_n^{(m)}), \leq_R)$
- every $p \in \mathcal{D}_n^{(m)}$ can be written as a join of elements in $\mathcal{J}(\mathcal{T}_n^{(m)})$

A CONJECTURE

For $m, n > 0$ we conjecture $\mathcal{T}_n^{(m)} \cong \beta \circ \delta(\mathcal{T}_n^{(m)})$, where

- $\beta = \beta_{m-1,m} \circ \dots \circ \beta_{2,3} \circ \beta_{1,m} \circ \dots \circ \beta_{1,2}$ and
- $\beta_{i,j} : (\mathcal{D}_n)^m \rightarrow (\mathcal{D}_n)^m$ is defined by replacing in (q_1, \dots, q_m) the i -th and j -th component by $q_i \wedge_R q_j$ and $q_i \vee_R q_j$, respectively

THEOREM: TAMARI-LIKE LATTICES FOR DIHEDRAL GROUPS

For $k > 1$ let \mathcal{C}_k denote the bounded poset whose proper part is the cardinal sum of a $(k-1)$ -chain and an atom.

Then for every $m > 0$, the poset $\mathcal{C}_k^{(m)}$ is a trim lattice, and its cardinality is $\text{Cat}^{(m)}(\mathfrak{D}_k) = \binom{m+1}{2}k + m + 1$.

EXAMPLE: THE LATTICE $\mathbf{DM}(\mathcal{T}_4^{(2)}) \cong \mathcal{T}_4^{(2)}$

